

Building A Secure Model of e-Medication Management System for Optimum Pharmaceuticals Safety

Syful Islam ^{#1}, Ratnadip Kuri ^{#2}, Md. javed Hossain ^{#2}, Dr. Humayun Kabir, ^{#3}
¹ syfulcste@gmail.com, ² ratnadipkuri@gmail.com ² javed.nstu@gmail.com ³ hkabar269@gmail.com

[#] Department of Computer Science and Telecommunication Engineering (CSTE), Noakhali Science and Technology University Sonapur, Noakhali-3814, Bangladesh

Abstract

Managing the processes of prescribing, dispensing, administering and monitoring of medicines is a fundamental part of ensuring good health outcomes and misuse use of medicines. Although improving medications management in Bangladesh is a key objective of the National drug policy-2016, there is no practical administration system for effective regulatory control. Our proposed e-Medication Management system (eMM) seeks to support policies and objectives by improving the access and quality of medicines information through the use of digital health service (eMM), enabling consumers and healthcare providers to make safe, informed decisions and achieve better health outcomes. We also investigate how eMM system relates to significant improvement of public health service.

Keywords:

eMM, Security, National Drug policy, Public health, Bangladesh

untreat- able infectious disease, so it is very much important to find out the reasons behind resistance and take some necessary steps to solve the problems. E-Medication management (eMM) is intended to prevent harm arising from ineffective medication information as patient's transition across healthcare settings. Although it is very critical component of providing safe care, it is complex and challenging to implement such system because the success of developing and integrating technical solutions to support eMM is highly dependent on attention to implementation processes, extensive usability testing and evaluation involving clinicians and other end-users. The main objective of this research is to develop an effective eMedi- cation Management system to control antibiotic drug selling in Bangladesh which will be transparent, accurate and ensure quality with efficacy. In addition, the eMM system will be implemented in such a way so that it can be easily accessed by general user from smart devices while ensuring all security perspectives.

I. INTRODUCTION

Medication is an essential part of health care and the appropriate treatment with drugs enables the cure and prevention of many conditions [1]. However, drug-related problems (DRP) are frequent and cause suffering for patients, and substantial costs for society [2], [3], [4]. A good medicines management can help to reduce the likelihood of medication errors and hence patient harm. Currently there is no standard for medication management in Bangladesh although antibiotic drug law 2016 is passed in cabinet. Among the risks for a country of the ineffective management of antibiotic drug supply are: introduction of drugs of unknown quality, unlabeled or expired drugs, drugs withdrawn from sale in other countries and being dumped or drugs produced cheaply without the necessary regard for quality; inadequate labeling of drugs (a particular problem where many people are illiterate), so that users are not properly informed on dosage, expiry date. Since uncontrolled antibiotic drug sales in our country causes a catastrophic increase in the medical and socio economic burden of

II. BACKGROUND

Self-medication is the use of drugs to treat self-diagnosed disorder or symptoms, or the intermittent or continued use of a prescribed drug for chronic or recurrent disease or symptoms [5]. In Bangladesh, it is very common that patient diagnoses his own illness and buys a specific drug to treat it. Besides, self-medication with antimicrobial compounds is an extremely ubiquitous practice in developing countries like us for reasons of cultural tradition, convenience, accessibility, perceived savings and other benefits, with poor adherence being a never-present consequence of self-medication. The arsenal of antibiotics may be used up quickly, leaving the patient vulnerable to drug resistant infections. Treatment of bacterial infections in the hospital and community has

been altered drastically over the past few decades with the emergence of pathogenic organisms that are no longer susceptible to most commonly prescribed antibiotics. There is a clear relationship between the amount of a given antibiotic used and the incidence of bacterial resistance. Resistance to commonly used antimicrobial drugs is remarkably high in our country

TABLE I
SUMMARY OF THE OUTCOME OF ANTIBIOTIC DRUG
MISUSE IN
BANGLADESH

Pub.	Study type	Purpose	Result
[6]	Survey	Assess and evaluate the prescription pattern of antibiotic	Polypharmacy
[7]	Survey	The amount of antibiotic prescribed by physicians	Irrational medication
[8]	Survey	The prevalence of self-medication	Self-medication
[9]	Survey	Situation about antimicrobial resistance	Lack of proper regulation
[10]	Case study	Role of pharmacist	Need of pharmacist to counsel

medical and socio economic burden of untreatable infectious diseases.

Moreover, over-prescribing and inappropriate prescribing are very common due to unethical practice of both health professionals and drug manufacturers. In addition, antibiotic self-medication is highly prevalent in here due to easy availability and poor regulatory controls on distribution of these drugs. Besides, patients can get antibiotics easily from pharmacies according to their demand without prescription. So, misuse of antibiotics is increasing. In long term effect, patients are exposed to serious health hazards, particularly liver and kidney damage, because of indiscriminate taking of antibiotics. Experts states that, unnecessary use of these drugs builds antibiotic resistance in human bodies, multiplying the extent of diseases that can even lead to death in extreme cases. In addition, if the resistance continues to develop, bacterial infections could go beyond treatment in future. Dr. Iqbal Arslan, professor of Biochemistry at Bangabandhu Sheikh Mujib Medical University(BSMMU) said, "Irrational use of antibiotics poses a great danger, as it develops drug resistance faster than what a new generation drug takes to be developed". He also blamed it on the counter sales of antibiotics, suggesting the government frame a strict law to prevent sales of antibiotics and painkillers without prescriptions of registered doctors. In table I we summarize the outcome

since antibiotics are not restricted. Increasing rates of antimicrobial resistance have left clinicians with limited drug options for the treatment of bacterial infectious diseases which is a major public health concern worldwide. Antibiotic resistance in developing countries causes a catastrophic increase in the of antibiotic drug misuse from paper [6], [7], [8], [9], [10] in Bangladesh.

III. LITERATURE REVIEW

Electronic medication systems are a crucial cornerstone of digital health system, because prescribing and dispensing medication is a frequent everyday activity that links various actors in the system (See [11]). Previous research has identified potential health and social benefits of e-prescriptions, such as improving quality of care and patient safety through error reduction(see [12]), and increasing patient convenience, particularly for repeat prescriptions(see [13], [14]). This paper reviews the existing literature on e-prescriptions/e-medication, focusing on the country (Estonia, United Kingdom, Sweden, and Denmark) case studies, and systematizes the evidence around benefits and success factors.

In Estonia, 1.3 million inhabitants, is known for its cutting-edge approach to digitization. In line with many European countries, the Estonian health system is founded on solidarity, with a single public payer, the Estonian Health Insurance Fund (EHIF), providing mandatory health insurance for nearly all of the population (See World Bank Group, 2015). The Estonian e-medication system enables data exchange between patients, providers, pharmacies and EHIF. To issue a prescription, the provider creates an entry in a patient's shared medication record, based on which patients can obtain their medication in any pharmacy in the country based on their eID. Patients can also log in via an online portal and view the audit trail of data access and use. Since the digital service has grown rapidly: 84 percent of prescriptions in the country were issued electronically in 2011, 95 percent in 2013, and over 99 percent today. According to a survey "Citizens' satisfaction with health and healthcare" 92% of users of digital prescription are satisfied with the service (see [15], [16]).

In United Kingdom, for its 64.1 million inhabitants, has a National Health Service (NHS) financed through general taxation and free at the point of care, which is separately managed for its constituent countries (England, Wales, Scotland, and

Northern Ireland). Digital services in health and other areas of government form part of the UK government's "digital by default" vision to increase service levels and reduce costs in the long term (see [17]). In the UK, large differences for e-prescriptions exist between primary and secondary care(see [18], [19]). In primary care, e-prescriptions are becoming relatively well established. In secondary care, e-prescription systems are largely implemented in a decentralized manner based on initiatives of specific hospitals (see [20]). The aim of the UK is to digitize hospitals and other aspects of the health system fully by 2020(see [21]).

In Sweden, with a population of about 9.5 million people, has a long history of the use of information technology in healthcare. It became one of the first countries to use e-prescriptions when, in 1983, a few doctors connected to local pharmacies to exchange prescriptions (see [22]). The national effort to improve connectivity in the health system began in 2000, when of health information technology and may be seen among the leading countries(see [25]). Danish citizens can use an online portal to access their health data and communicate with health providers. E-prescriptions began in Denmark in 1994, with the central eHealth organization Medcom driving stakeholder alignment and setting up the necessary standards for e-prescriptions. In 2010, about 88 percent of all prescriptions were sent electronically in Denmark; as of 2016, there are virtually no paper-based prescriptions anymore.

In Spain, The electronic transmission of prescriptions to pharmacies is a regional routine. There is an uneven implementation of electronic prescription within Spain and interoperability of the regional e-prescription systems is an issue. Andalusia has an advanced system than the other Spanish regions. Regions with e-prescription systems reported the following e-prescription take up during 2008–2009: Malaga 56.3 %, Balearic Islands 53.46 %, Cordoba 50.34%, Extremadura 50.3 %, Andalusia 46 %, Almeria 41 %, Mallorca 14.12%, Menorca 3.04 %. Madrid is undergoing a pilot and Navarra, Cantabria, Murcia, Ceuta and Melilla are preparing pilots (see [26]).

In Australia, health system is world class, supporting universal and affordable access to high quality medical, pharmaceutical and hospital services, while helping people to stay healthy through health promotion and disease prevention activities (see [27]). Electronic prescription in Australia is currently provided by two service providers, MediSecure and eRx. Both services can be integrated into many of the existing clinical and pharmacy prescribing software systems. Since December

2012, they have become interoperable allowing bilateral transfer of information. In Canada, On March 22, 2016, the Government of Canada allocated funds to Canada Health Infoway to develop an e-prescribing service. Infoway is working with Health Canada, the provinces and territories and industry stakeholders to create PrescribeIT, a multi-jurisdiction e-prescribing service. In USA, In the United States, the HITECH Act promotes adoption of this technology by defining e-prescribing as one meaningful use of an electronic medical record. According to data released in May 2012 by Superscripts, a company which operates the nation's largest health information (e-prescribing) network, roughly 317,000 office-based physicians now e-prescribe in the United States. A more recent report released by the Office of the National Coordinator for Health IT in June 2012 finds that 48 percent of U.S. physicians use e-prescribing systems (see [28]). National growth in e-prescribing over the period September 2008 through June 2012 increased over 40 percent, with individual states increasing adoption anywhere from 28 percent to 70 percent (see [29]). In Bangladesh, Misuse of antibiotic medicine is at alarming state. Over prescribing, inappropriate prescribing by doctors, over counter availability of drugs are very common here. In order to reduce misuse of drugs, the government of Bangladesh has approved the draft of revised and integrated drug policy, first of its kind in Bangladesh, to make sure pharmaceutical products meet international standards and substandard, fake and adulterated drugs are checked. The National Drug

2012, they have become interoperable allowing bilateral transfer of information. In Canada, On March 22, 2016, the Government of Canada allocated funds to Canada Health Infoway to develop an e-prescribing service. Infoway is working with Health Canada, the provinces and territories and industry stakeholders to create PrescribeIT, a multi-jurisdiction e-prescribing service.

In USA, In the United States, the HITECH Act promotes adoption of this technology by defining e-prescribing as one meaningful use of an electronic medical record. According to data released in May 2012 by Superscripts, a company which operates the nation's largest health information (e-prescribing) network, roughly 317,000 office-based physicians now e-prescribe in the United States. A more recent report released by the Office of the National Coordinator for Health IT in June 2012 finds that 48 percent of U.S. physicians use e-prescribing systems (see [28]). National growth in e-prescribing over the period September 2008 through June 2012 increased over 40 percent, with individual states increasing adoption anywhere from 28 percent to 70 percent (see [29]). In Bangladesh, Misuse of antibiotic medicine is at alarming state. Over prescribing, inappropriate prescribing by doctors, over counter availability of drugs are very common here. In order to reduce misuse of drugs, the government of Bangladesh has approved the draft of revised and integrated drug policy, first of its kind in Bangladesh, to make sure pharmaceutical products meet international standards and substandard, fake and adulterated drugs are checked. The National Drug

Policy 2016 will replace the existing one framed in 2005 (see [30]). According to the policy, no one can buy drugs without prescriptions from physicians, except 39 allopathic, 23 ayurvedic and 48 unani medicines. This is very important as indiscriminate use of medicines, especially antibiotics, is harmful for public health. The policy also provides for availability of effective, safe and standard drugs, rational and safe use of drugs, registration of drugs, registration for importing drugs as well as manufacturing drugs and its raw materials. There are proposals as well for the formation of a national drug regulatory authority, advertisement control, publicity of drugs and joint initiatives for research and development. The revised policy makes storage or display of drugs with the expiry dates changed or distorted a punishable offence. Legal steps would be taken against people for charging customers additional prices. Once the revised policy is implemented, there will be little scope for sub-standard and adulterated drug manufacturers (see [30]). But there is no e-medication framework (Table I) to support the policies and objectives to improve current situation in Bangladesh. Now it's time being demanding higher standards of care, better patient safety, more value for money spent, and an integrated continuum of care delivery.

IV. ELECTRONIC MEDICATION MANAGEMENT SYSTEM

Medication management (MM) is a formal process in which healthcare providers work together with patients and families to ensure accurate and The data collected was subjected to simple descriptive statistical analyses including frequency distribution and percentage, since the purpose of the study was to find irrational antibiotic medication pattern and propose a framework to ensure fair antibiotic medication. From Patients survey, it is marked that 40% take antibiotic medicine by taking prescription from doctors and 60% take antibiotic medicine without taking prescription from doctors due to suffering cough and fever most of the time. Besides, from doctors survey, it is evident that 55% prescribe antibiotic medicine in suspected cases and 45%

comprehensive medication information communicated consistently a cross transitions of care. The goal of MM is to prevent potential patient medication errors and adverse drug events. So in formal definition, Electronic Medication Management System (eMM) uses information technology to access and integrate electronically stored patient medication data to support: (i) the collection of the electronic Best Possible Medication History (eBPMH), the detection and resolution of discrepancies, (ii) the development of an electronic Best Possible Medication Discharge Plan (eBP- MDP) and (iii) the comparison of eBPMH and new orders at transfers. Moreover, eMM system can be tailored according to the policy and objectives of a specific country and organization (see [12]).

V. SURVEY ON THE NECESSITY OF EMM SYSTEM

Generally success of e-administration depends on how users of the system accept it as a service. Implementation of an eMM will require changes in workflow. Since e-Medication Management system is completely new idea in perspective of Bangladesh, it is important to understand the services required, degree of acceptance, prospect and challenges to implement the system. In order to understand the necessity of e-medication system we run a paper based end-to-end survey that cover three basic entity of the system (i.e. doctors, patients and Medicine sellers). Our questionnaires were given only to the spontaneously interested candidates during the survey.

prescribe medicine after confirming disease which is a clear indication of irrational medication. Most of the doctors, patients, sellers opine that in order to avoid serious health hazard, patients should stop antibiotic drug self-medication, and they also blame that over counter availability of antibiotic drug is also a major reason of irrational medication (See figure 1). At the end of the survey, we observed that most of them (patients, doctors, medicine sellers) feel to use an antibiotic e-administration system that will ensure fair antibiotic medication practice (See figure 2, 3, 4).




Fig. 1. Illustrating opinion on “misuse of antibiotic for over counter availability of antibiotic”.




Fig. 3. Illustrating opinion on “None get medicine without e-prescription”.




Fig. 2. Illustrating opinion on “Should we use e-medication system”.




Fig. 4. Illustrating opinion on “e-Medication system can improve public health”.

VI. PROPOSED SYSTEM MODEL

The main objective of this paper is to design and develop a drug Administration framework that will help ministry of health effectively in regulatory controls on the distribution

and selling of antibiotics in order to reduce the frequency of antibiotic misuse. Since e-Medication system is a complex system, it must include all service and security component. So Prior to our design and implementation, we accomplished our requirements engineering processes by interacting with the doctors, patients, medicine sellers to gather the requirements for the system (See figure 5). We were able to identify some services which are represented architecturally in the following sections:

A. SERVICE REQUIREMENTS

The IT infrastructure for medication management should support all the clinical and administration services involved and should include the following:

- Comprehensive medication profile: The core component of a medication management infrastructure is a comprehensive medication profile that captures information about all

medication transactions. More specifically, a comprehensive medication profile will contain: a history of all previous dispensing, a history of all non-dispensing events (such as “stop orders”), an




Fig. 5. Simplified presentation of e-Medication system Model.

reduce medication errors, support appropriate, prescribing practices, and streamline medication management interactions between physicians and pharmacists. Most electronic prescribing systems function in the following way: The physician sends a prescription to an approved repository, allowing a pharmacy to retrieve the prescription with the patient’s authorization. Pharmacies retrieve the electronic prescription from the approved repository using a unique identifier. If the physician cannot send the prescription electronically to the pharmacy selected by the patient, the physician prints out a hard copy of the prescription for the patient (A.B.R. Kumar, et al, 2012).

- knowledge management and translation: to provide appropriate, 24-hour clinical and drug information’s.
- Provider and patient portals: to give patients and their health care providers ready access to patient information and to ensure greater involvement of patients in their own health care. Provider portals make it possible for health care providers to get access to information and resources specific to their practice. For example, electronic health journals and news, Practice guideline, secure email etc.
- Pharmacy network: Advanced medication management systems depend on secure, broadband networks integrated with other health infrastructure at the local, regional, and provincial levels. A pharmacy network must comply with standards for connectivity and compatibility (such as the National claims Standard Initiative) to support pharmacy

active prescription drug profile; information on past drug interactions.

- Electronic prescribing: Electronic prescribing systems to claims processing and adjudication by public and private health insurance companies.
- Standards: to ensure that the entire system is integrated and coordinated;
- Pharmacy management system: to support dispensing, in-ventory management, claims processing and adjudication, and compensation;
- Medical test report management system: to support up-loading the lab test report in the specific patients account so that doctors can view these report according to their demand.
- Physician practice management system: to manage the

administrative functions of a physician office, such as appointment scheduling and physician billing.

B. SYSTEM REQUIREMENTS

In order to make completely successful and long lasting system, it must fulfill the following requirements. It includes:

- System reliability: System reliability is an essential aspect of implementing eMM. There must be a back-up plan if eMM fails or goes down.
- Usability: Higher rates of compliance will be achieved with an intuitive, easy-to-use system. Users are more likely to revert to paper-based processes or find workarounds if eMM is difficult to use.
- Sustainability: To ensure sustainability, it is important to continually consider and communicate: eMM achievements and remaining goals, the importance of eMM from a patient safety perspective, Buy-in and support from clinicians and other end users of the system.
- Cost: For any implementation to be a success it is important to identify project and long term maintenance costs. Therefore it is critical: Ensure commitment and support of senior management through an executive sponsor dedicated to this project, Ensure that sufficient financial resources are available to implement eMM, Provide a dedicated team of healthcare professionals and Health IT staff for project implementation, Determine the impact on time of using new eMM processes by physicians, nurses and pharmacists, Ensure there is sufficient staff with enough time to conduct eMM on an ongoing basis.
- Patient Safety: As with other electronic tools, the introduction of eMM has the potential to reduce errors and/or introduce new errors into the eMM process. To fully benefit from the potential of eMM to decrease errors: Ensure that healthcare professionals are educated in and aware of their role in all aspects of eMM processes, Encourage healthcare professionals and patients to report near misses and errors that arise, in order to refine eMM processes to ensure system reliability and maximize the benefits of eMM, Ensure a well-defined evaluation plan is in place to track compliance with and outcomes of implementing eMM.
- Risk Assessment: Although eMM has many benefits, implementation may introduce risks and potential for error that must be considered and mitigated. Risks potentially introduced by eMM implementation include: Over-reliance on electronic medication lists and avoidance of, or performing lower quality, patient interviews. Technology-induced errors may be inadvertently introduced and inadvertently increasing workload by requiring electronic entry of medications.
- Training: Training of users is critical and clinicians will require more than one training session or training combined with follow-up support. Indeed, in studies of eMM implementations, training and education of end users of eMM was found to be one the most important factors in facilitating adoption by physicians and pharmacists

VII. ARCHITECTURE OF THE SYSTEM

Since the e-Medication system is an emergency service portal it needs to be characterized by timely and accurate access of information at all time. We design an architecture that will always allow easy access, search, retrieval and update of information from access device. In our system, we use client server multi-tier based architecture for hardware (physical) and 3-tier client/server architecture for software (logical) deployment.

A. HARDWARE ARCHITECTURE

Our system architecture comprises of a complete range of robust performance client and server platforms with

integrated enterprise application as shown in figure 6. The hardware architecture is highly secure and utilizes multiple layers of firewall protection to create several region of trust. The robust servers with load balancing system provides, real-time access to point of care database originating from systems across the enterprise system to facilitate timely and accurate care delivery and practice management. From client side, information can be accessed through PC, smart phone and other internet enabled devices. User Load on the application server tier is balanced by using multiple application servers and load balancer. Firewall is integrated to filter all traffic moving in and out of the system. When a user proxy database is created, metadata for the proxy tables is imported automatically from the remote location that contains the actual tables. This metadata is then used to create proxy tables within the proxy database to make more secured database back-up.




Fig. 6. Hardware architecture for e-Medication Management (eMM) system.

B. SOFTWARE ARCHITECTURE

Figure 7 gives an overall logical (software) view of the architecture of the eMM system that shows the locations of the each of the supports service in the system. Our system is a 3-tiered client-server which consists of the client interface, middleware and database repository where database is separated from the client through the middleware. The middleware concept helps to solve scalability, load balancing, transactional processing and interoperability issues by providing a means that allows different hardware and software from different manufactures to share common Patient medical records. The client has its parts the Security and Authentication Support Service (SASS), Documentation Support Service (DSS) and




Fig. 7. Logical architecture of e-medication management system.

the proper identity or not. If the user account contains proper identity, the user will be proceeded to use the system. If not, the user account will be pended to use and detected as a malicious user. Then the system will

the View and Report Service (VRS). These support services do not store or process any form of data. They only provide an interface for middle layer and the data layer.

C. HOW IT WORKS?

The system is based on forcing the user except patient to register the real identity to ensure transparency in medical practice. This System has Databases, managed by the system authority as well as Database Manager. The whole system (see figure 8) comprises of two types of users: 1) Who use the system 2) Database Management Authority. Through this process eMM system will help fair medical practice.

D. SOFTWARE ARCHITECTURE




Fig. 8. Diagram of prescription flow and drug dispensing cycle.

1) USER OF THE SYSTEM: In our system, there are three types of user. They are medical practitioners (i.e. doctor, Medical technicians), patient and seller. In our system, Patients registration process is optional keeping in mind about emergency medical service but doctor and medicine seller must be registered user to interact with the whole process of prescription service. Our System will force the users to use real identifications. There will be verification technique in our system to check whether the user account holder use

automatically notify them to use proper identity to prevent from being suspended. After completing the registration process successfully doctor can view medical history (i.e. disease history, previous pre-

scriptions etc.) and prescribe medicine to the patient through our drug administration system. When a prescriber (doctor, specialist, dentist etc.) generates an e-Prescription two things happen - an electronic copy of the prescription is secured via encryption and submitted to the Prescription Exchange Service (PES); and a unique number is generated and printed on the paper prescription. When the patient takes the paper prescription to the participating pharmacy, pharmacists or authorized sellers will distribute drug to the respective patient. In addition, the system will automatically send SMS to the prescription holder, if medicine is taken by another person and track basic information of the buyer. If the pharmacy does not use eMM system, the prescription is dispensed as per the standard dispensing process. That is, the prescription information is manually transcribed into the pharmacy's dispensing system and the medicine dispensed. Moreover, system will contain facility of partial dose purchasing facility. Besides it will also contain feedback facility so that doctors and patients can communicate with each other to know whether the disease is cured or not. This system can be accessed using both computer and smart phone for ease of use.

2) *DATABASE REPORTING AUTHORITY*: In order to ensure fair practice, the system will be under proper authority. The authority will periodically check the user's activity. If a user of the system want to complain about any illegal activity then he/she will enter into our proposed system and paste that id of illegal practitioner in the mentioned box. Then he has to write a description why he complains against the ID. Then he has to click the report button. Finally the reported complains will be saved in the system Database. Here one thing should be mentioned that only registered and valid ID holder can report their complains in our system model.

E. ANTIBIOTIC DRUG REGISTRATION AND CATEGORIZATION: Authorizations to manufacture, import, distribute or sell antibiotic drugs in Bangladesh are only being granted for registered antibiotic drugs which are included in the National health database. In our system, there will be an updated list of antibiotic drug authorized by ministry of health Bangladesh. Though our system only registered antibiotic drug can be prescribed and sold. For special reasons and after receiving opinion of the health ministry, the Minister may authorize the distribution of un-registered antibiotic drugs not yet included in the database. Our system contains the facility to update the antibiotic drug list.

F. SECURITY DESIGN

In the e-Medication Management (eMM) system scenario each asset has huge importance as long as the system must TABLE II

SUMMARY OF ROLE BASED ACCESS CONTROL MECHANISM FOR USERS OF THE SYSTEM

Entity	Read Permission (Patients Personal information + Medical history + Registered drug list+ Dose Information)	Write permission (Patients Personal Information + Medical history + Registered drug list+ Dose Information)
Patient	Yes (All)	Yes (Patients Personal Information)+ No (Medical history + Registered drug list+ Dose Information)
Doctor	Yes (All)	Yes (New Medical History+ Dose information)+ No (Patients Personal Information+Registered drug list)
Medical Technician	Yes (Patients Personal Information+ Registered drug list+ medical history)+ No(Dose Information)	Yes (Medical Lab Report)+ No (Patients Personal information +Medical history+Registered drug list+ Dose Information)
Drug seller	Yes (Dose Information+ Registered drug list+ Personal Information)+ No (Medical History)	Yes (Dose Information)+ No (Patients personal Information+Medical history +Registered drug list)
System administrator	Yes (All)	Yes (Patients personal information)+No(Medical history+ Registered drug list)

provide services with high reliability and guarantee the confidentiality of the data. We have secured eMM system by applying security mechanism I) Availability II) Authentication and Access control III) Confidentiality IV) Non-repudiation at different layers.

- *Availability*: Since this system is an emergency service portal it must be available to the users at all time from everywhere. System availability is ensured by robust set of servers. Besides load balancer technology is integrated to ensure availability of service as per user request. In addition firewall is implemented to filter out unusual traffic in and out from the system.
 - *Authentication and Access Control*: In order to provide authentication and access control we divide this scenario into two cases. General user case: Our System will force the

users to use real identifications. There is proper verification technique using National ID, Birth registration number, Driver license number, passport number for general user, BMDC number for doctors and drug license number for medicine sellers to check whether the user account holder use the proper identity or not. If the user account contains proper and doctors activity related to medical service seems unusual in their account then they can complain against them (doctor and patient vice versa) in our system. Besides access control is another security approach since our system database contains sensitive data of users. In our system we use role based access control mechanism for users as shown in table II

- Emergency User case: In case of emergency use, if a user has no account he/she can take medical service using his any unique ID (Ex: national ID, Birth certificate number, Passport number, driver license, student ID etc.). In this case the doctor will generate a prescription ID for the patient to prescribe medicine and print a prescription with a unique ID. Using this prescription ID, the patient/ relative of the patient can take medicine from registered medicine store. In order to handle misuse of the emergency service, the system will contain facility to check frequency of taking medicine by the patient from different doctor in the same day using same ID. Using emergency medical service, a patient can take medicine from doctors not more than n ($n=2$ or 3 or 4) times for same purpose. If he/she requires medicine further, he/she must have to register as valid user of our system through proper registration process and must require special authorization from doctors.
- Confidentiality: The major documents that are to be exchanged among totally different medical entities are patient's medical reports. These documents ought to be changed firmly. In our system all reports of the patient are encrypted before uploading to the patient's medical directory. To look at an improved security performance a cryptologic rule will be used. Therefore, a trade-off between security and system performance has to be thought about fastidiously. Therefore AES encoding rule can be employed that is optimum (Key size=128, 192, 256 bit, Block

identity, the user will be proceeded to use the system. If not, the user account will be pending to use and detected as a malicious user. Then the system will automatically notify them to use proper identity to prevent from being suspended. Thus fake account holder will be filtered and reduce misuse of our system. If a patient

Size=128 bit, speed=Fast (use computer memory unit substitution, row shifting, column compounding key auditing), Rounds=10, 12 or 16, crackable=Not possible) (A.B.R. Kumar, 2012). In our planned system we tend to use AES Rijndael-128 algorithm (key size=128) to cipher and rewrite a file. To implement file encoding and cryptography method we've got used PHP Avestan Framework one.11.0.

- Non-repudiation: Most of the cases suspected medical employees are unidentified because of the non-transparency of the system. Medical employee's offers health service through eMM system and a central access log is maintained to trace every medical entity. Therefore suspected medical employees are going to be known at the time of event or straight off when the event through this access log.

G. SYSTEM IMPLEMENTATION AND UNIT TESTING

The proposed model has been developed at the Computer Lab of NSTU and tested in lab environment by dividing students into three categories of actors (i.e. Doctors, patients and medicine sellers). Teachers and Students have played the role of patient, doctor and seller. We assume our testing environment has the facility of patient treatment, diagnostic test and pharmacy. The testing of this model has been applied on 40 users (students, teachers). The web based module with database server has been placed at NSTU data center. Testers of the system has played the role of central admin to verify register users because currently there is no instant access on national, BMDC and pharmacist databases for verification of patients, doctors and medicine sellers. Figure: 9 shows the main page (web module) of the e-health system where central admin is logging to system to perform his task. Each medical entity can login and can perform their task through this page.




Fig. 9. eMMS system demo page view.

VIII. BENEFITS OF EDUCATION MANAGEMENT SYSTEM

Better medication management means that the health system and health professionals are organized to ensure that the most appropriate medications are prescribed and consistently taken by the patient with desired results and fewer adverse reactions (Emergis Inc., 2006). The implementation of eMM system in Bangladesh improves the accuracy of the medication related functions, transparent with the legibility issues related to a tradition medication flow. By using an eMM system users get benefits included:

with doctors for further suggestions and feedback. Moreover, it will highly reduce irrational medication problems and consequent result is improved public health.

- Benefits to medicine sellers: eMM system has the potential to significantly improve pharmacy dispensing operations. Prescriptions that are electronically transmitted are more legible and can improve work flow. More effective handling of refill authorization and routine issues via electronic means will also improve the efficiency of pharmacy operations.

XI. CHALLENGES TO IMPLEMENT AN E-MM SYSTEM

In order to implement this eMM system in national level, we have to overcome some limitations. These are:

- Financial Cost and Return on Investment (ROI) - The costs related with purchasing, implementing, supporting and maintaining such a system may be beyond the means of most small clinical practices in

- Benefits to Doctors: When eMM system is fully

implemented in the office environment, Doctors find that efficiency increases. eMMs applications have the potential to check drug information databases for appropriate prescribing guidelines, the patient's complete medication profile for drug interactions, and the patient's electronic medical record for disease contraindications. These features save the doctors valuable time and effort and further reduce the number of errors or adverse events.

- Benefits to Patients: The benefit most evident to patients is increased convenience. Prescription orders sent via electronically to pharmacies make it possible for patients to arrive at the pharmacy and have their prescription orders waiting for them. Patients can easily communicate specially in rural area, and which we noted it to be one of the greatest implementation barriers.
- Change of management – Healthcare providers and associated staff are adapted their current management system, in which case change of management becomes extremely important. Also the management needs to be trained to adapt with new system.
- Integrity of data input - Accidental data entry errors such as selecting the wrong patient or clicking on the wrong choice in a menu of dosages may occur. In this case Software vendors can reduce errors by continually reviewing user feedback and follow best practices in user interface design.
- System downtime - Periods of system downtime may arise, either due to network-related issues, hardware failure, or loss of electricity. In addition, the inability to use electronic prescribing when the system is not accessible is of great concern, and must be addressed with the discussion of fall-back procedures and mechanisms when such situations arise.
- Patient Access Lost - In the event of a development

beyond the control of the patient, such as a software malfunction in the health care provider's office, the patient can no longer ask the care provider for a paper script to take to a pharmacy in order to obtain needed medicines. This leaves the patient at the mercy of technicians or other unrecoverable workers.

XII. COMPARISON OF eMM SYSTEM WITH TRADITIONAL PAPER BASED SYSTEM

The following comparison (Table III) will show how eMM system significantly improve public health service in Bangladesh keeping consistency with National drug policy 2016.

TABLE III COMPARISON OF eMM SYSTEM WITH TRADITIONAL PAPER BASED MEDICATION SYSTEM

Parameters	eMM System	Traditional paper based system
Reduce prescribing and dispensing errors?	Yes	No
Improved standardization of documentation?	Yes	No
Improved accessibility of documentation?	Yes	No
Avoid more adverse drug interactions and reactions?	Yes	No
Potentially improved compliance through the implementation of soft stops?	Yes	No
Improved efficiency of many medication-related processes?	Yes	No
Reduce the incidence of drug diversion?	Yes	No

- [3] A. J. Avery, A. Sheikh, B. Hurwitz, L. Smeaton, Y.-F. Chen, R. Howard, J. Cantrill, and S. Royal, "Safer medicines management in primary care." *Br J Gen Pract*, vol. 52, no. Suppl, pp. S17–S22, 2002.
- [4] F. Salvi, A. Marchetti, F. D'Angelo, M. Boemi, F. Lattanzio, and A. Cherubini, "Adverse drug events as a cause of hospitalization in older adults," *Drug safety*, vol. 35, no. 1, pp. 29–45, 2012.
- [5] W. H. Organization *et al.*, "Guidelines for the regulatory assessment of medicinal products for use in self-medication," Geneva: World Health Organization, Tech. Rep., 2000.
- [6] M. M. Begum, M. S. Uddin, M. S. Rahman, M. A. Nure, R. R. Saha, T. Begum, R. Begum, A. Islam, M. Sultana, and R. Karim, "Analysis of prescription pattern of antibiotic drugs on patients suffering from ENT infection within Dhaka metropolis, Bangladesh," *International Journal of Basic & Clinical Pharmacology*, vol. 6, no. 2, p. 257, 2017.
- [7] M. A. Sayeed, N. Iqbal, M. S. Ali, M. M. Rahman, M. R.

XIII. CONCLUSION

The current method of prescribing, dispensing, administering and monitoring for medication is completely ineffective and under tremendous pressure to change. Bangladeshis are demanding higher standards of care, better patient safety, more value for money spent, and an integrated continuum of care delivery. Our proposed electronic medication management (eMM) system with information technology can help to achieve these goals. We have developed a centralized eMM system with all medical entities (Doctors, Patients and Medicine Sellers) to give faster service to the patient by replacing paper based medical record by web based medical record (e- Prescription) where the privacy of all entities has been ensured by role based access. Malpractice by doctors, over-counter drug availability, and irrational medications can be completely controlled through our system. Also through this project implementation, a strong nexus among patients, physicians and drug sellers will be established which will help fair exercise of digital medical practice.

ACKNOWLEDGMENT

The research group would like to thank doctors, medicine sellers, students who participated in the survey of this research work.

REFERENCES

- [1] E. Topinková, J. P. Baeyens, J.-P. Michel, and P.-O. Lang, "Evidence-based strategies for the optimization of pharmacotherapy in older people," *Drugs & aging*, vol. 29, no. 6, pp. 477–494, 2012.
- [2] P. M. van den Bemt, T. C. Egberts, J. R. Brouwers *et al.*, "Drug-related problems in hospitalised patients," *Drug safety*, vol. 22, no. 4, pp. 321–333, 2000.
- Islam, and M. Jakaria, "Survey on antibiotic practices in Chittagong city of Bangladesh." *Bangladesh Pharmaceutical Journal*, vol. 18, no. 2, pp. 174–178, 2015.
- [8] M. Biswas, D. N. Roy, A. Tajmim, S. S. Rajib, M. Hossain, F. Farzana, and N. Yasmen, "Prescription antibiotics for outpatients in Bangladesh: a cross-sectional health survey conducted in three cities," *Annals of clinical microbiology and antimicrobials*, vol. 13, no. 1, p. 15, 2014.
- [9] M. S. Rahman and S. Huda, "Antimicrobial resistance and related issues: An overview of Bangladesh situation," *Bangladesh Journal of Pharmacology*, vol. 9, no. 2, pp. 218–224, 2014.
- [10] D. Saha, "Health care system in Bangladesh," *Journal of Pharmaceutical Care*, pp. 77–78, 2013.
- [11] R. Hibberd, N. Barber, T. Cornford, and V. Lichtner, "The evaluation of the electronic prescription service in primary care: interim report on the findings from the evaluation in early implementer sites," 2012.
- [12] A. Agrawal, "Medication errors: prevention using information

- technology systems," *British journal of clinical pharmacology*, vol. 67, no. 6, pp. 681–686, 2009.
- [13] T. Porteous, C. Bond, R. Robertson, P. Hannaford, and E. Reiter, "Electronic transfer of prescription-related information: comparing views of patients, general practitioners, and pharmacists." *Br J Gen Pract*, vol. 53, no. 488, pp. 204–209, 2003.
- [14] T. Hammar, S. Nyström, G. Petersson, B. Åstrand, and T. Rydberg, "Patients satisfied with e-prescribing in sweden: a survey of a nationwide implementation," *Journal of pharmaceutical health services research*, vol. 2, no. 2, pp. 97–105, 2011.
- [15] A. M. Aguilar Rivera and K. Vassil, *Estonia: A Successfully Integrated Population-Registration and Identity Management System*. World Bank, 2015.
- [16] A. Säär and A. Rull, "Technology transfer in the eu: exporting strategically important ict solutions to other eu member states." *Baltic Journal of European Studies*, vol. 5, no. 2, pp. 5–29, 2015.
- [17] N. Khilji, Y. Duan, R. Lewis, T. Bukoye, and U. Luton, "Incorporating knowledge management tools in the uk local government towards improved planning support services," in *ICICKM 2017 14th International Conference on Intellectual Capital Knowledge Management & Organisational Learning: ICICKM*, 2017, p. 122.
- [18] T. Cornford, B. Dean, I. Savage, N. Barber, and Y. Jani, "Electronic prescribing in hospitals-challenges and lessons learned." 2009.
- [19] S. Garfield, Y. Jani, S. Jheeta, and B. D. Franklin, "Impact of electronic prescribing on patient safety in hospitals: implications for the uk," *Acute pain*, vol. 10, p. 00, 2019.
- [20] G. C. Patton, S. M. Sawyer, J. S. Santelli, D. A. Ross, R. Afifi, N. B. Allen, M. Arora, P. Azzopardi, W. Baldwin, C. Bonell *et al.*, "Our future: a lancet commission on adolescent health and wellbeing," *The Lancet*, vol. 387, no. 10036, pp. 2423–2478, 2016.
- [21] N. Board, "Personalised health and care 2020—using data and technology to transform outcomes for patients and citizens—a framework for action," *London: HM Government*, 2014.
- [22] B. Åstrand, E. Montelius, G. Petersson, and A. Ekedahl, "Assessment of eprescription quality: an observational study at three mail-order pharmacies," *BMC medical informatics and decision making*, vol. 9, no. 1, p. 8, 2009.
- [23] G. O. Klein, "History of electronic prescriptions in sweden: from time-sharing systems via smartcards to edi," in *IFIP Conference on History of Nordic Computing*. Springer, 2010, pp. 65–73.
- [24] S. B. Care, "Eprescribing and electronic transfer of prescriptions: an international review," 2012.
- [25] P. Kierkegaard, "E-prescription across europe," *Health and Technology*, vol. 3, no. 3, pp. 205–219, 2013.
- [26] A. Corbellini, S. Giest, J. Artmann, J. Heywood, and J. Dumortier, "Country brief: Spain-health strategies," 2010.
- [27] M. L. Fleming and E. Parker, "Sample proofs@ elsevier australia."
- [28] M. D. Smith, "Abilify, brintellix, champix: The abcs of fda proprietary name disputes," *Cardozo Arts & Ent. LJ*, vol. 35, p. 805, 2016.
- [29] M. Hufstader, M. Swain, and M. F. Furukawa, "State variation in e-prescribing trends in the united states," *ONC Data Brief*, no. 4, 2012.
- [30] M. Islam *et al.*, "Factors behind exiting from the bangladesh pharmaceutical industry and closing down the pharma businesses unit of glaxosmithkline bangladesh ltd." 2018.